
2025

Professional Dairy Managers of Pennsylvania (PDMP)

Corn Silage Hybrid Performance
Trial Results



Prepared by: Alex Hristov (PSU Animal Sciences), Sergio Francisco (Cargill), Chris Canale (Cargill), Hanna Wells(PSU Plant Science), Dayton Spackman (PSU Plant Science), Cassidy Bumbaugh (PSU Plant Science), Charlie White (PSU Plant Science)

Production Details: PDMP Corn Silage Hybrid Evaluation Trials

Site:	West Grove, PA
Cooperator	Walmoore Holsteins Inc. (Walt Moore)
Planting Date	5/1/2025
Soil Type	Glenelg Silt Loam
Herbicides	pre- 1.2 qt Calisto Xtra, 1.2 qt Glystar 5 Extra (Roundup)
	post- 0.75 qt Calisto Xtra, 1 qt Glystar 5 Extra (Roundup)
Previous Crop	Corn followed by barley cover crop
Tillage	Fall cover crop turbo tilled in, corn no tilled into burned off cover
Starter Fertilizer	15 gal UAN
Insecticide	
Manure	18,000 gallons dairy liquid per acre. Per 1000 gallons - 10.71 pounds Ammonium Nitrogen, 7.62 pounds P2O5, 14.6 pounds K2O
Fertilizer pre plant	20 gal 27-0-0-3 with DuoMax Stabalizer
Fertlizer sidedress	None
Foliar extras	0.7 oz Source corn, 1 qt TimaUp, 1 qt Kfuel, 4 oz Nano Pro
Harvest Date	9/4/2025
Field Summary:	

Field Summary:

Despite the weather challenges this season posed, this location had even emergence and excellent yields. There was little to no animal damage and minimal disease present.

Weather Summary:		
Month	Precip.	GDD
May	6.7	363
June	1.7	655
July	7.8	860
August	2.5	639
September 1 - September 4	0.0	74
Seasonal Total	18.7	2591
Precip. Data:	http://wunder	ground.com
GDD data:	http://climates	martfarming

PDMP Corn Silage Hybrid Testing Program 2025

PROFESSIONAL DAIRY MANAGERS OF PENNSYLVANIA in collaboration with

I NOTES

PennState Extension
College of Agricultural Sciences

Late maturity (111-118) day RM silage hybrids in West Grove, PA

Notes: SEE BACKGROUND TAB
Cooperator: Walmoore Holsteins Inc.

					Dry	Crude						uNDF			Fresh		DOM	
		Relative	1	Pop.	Matter % ²	Protein	Lignin	Ash	Starch	TFA	NDFom	240 hr	NDFD 30	IVSD %Starch ³	Yield tons/ac⁴	OM Yield tons/ac⁵	Yield	OMD % ⁷
Brand	Hybrid	Maturity	Traits ¹	Plants/ac	%	%DM	%DM	%DM	%DM	%DM	%DM	%DM	%NDF	%Starcn	tons/ac	tons/ac	tons/ac ^b	%
111-114 day hybrids	XC24121	114	10	22.000	42 C	0.0	2.2	2.5	45.7	2.7	27.1	0.0	F2.0	67.4	20.0	٥٢	6.3	64.0
Advanta Seeds		== :	10	33,000	43.6	8.0		2.5	45.7	2.7		8.8	53.0	67.1	28.0	9.5	6.2	64.9
Dekalb	DKC111-02RIB	111	40	33,000	41.7	7.8	2.1	2.5	45.2	2.8	28.1	8.1	57.2	67.2	27.8	9.5	6.3	66.1
Growmark FS	INVISION FS 6349PC R/	112	32	33,000	41.6	8.0	2.0	2.4	47.3	2.9	25.2	7.6	55.4	67.4	29.6	10.1	6.7	66.2
Seed Consultants	SC1135PCE	113	33	33,000	41.4	7.8	2.0	2.4	45.9	2.7	27.1	7.8	56.8	67.5	25.4	8.7	5.8	66.3
Syngenta	E114C4-DV	114	19	33,000	41.3	7.7	2.0	2.4	45.2	2.9	27.6	8.4	55.5	67.4	27.7	9.5	6.2	65.7
Dekalb	DKC113-62RIB	113 113	40	33,000	41.0 40.9	7.8	2.0	2.5	45.6 46.5	2.8	27.3	7.7 7.7	57.3	67.8	29.3	10.0	6.7	66.5
Advanta Seeds	XC24921		19	33,000		7.8 7.7	2.0			2.8	27.1		55.9	67.7	28.8	9.9	6.5	66.1
Channel	214-95SSPRIB	114	40	33,000 33.000	40.8 40.6	7.7	2.0	2.6	45.9 44.1	2.8	27.1 29.7	8.4	54.6 55.8	67.3 67.0	25.8 27.5	8.8 9.4	5.8 6.1	65.3
Dekalb	DKC61-80RIB	111	39 40	,		7.5	1.8	2.5	44.1	2.7		9.0 7.6			27.5	9.4	6.6	65.3 66.7
Channel	213-53SSPRIB	113		33,000	40.6						28.9		60.1	67.0				
Seedway	SW 1331SP	113	42	33,000	40.6	7.8	2.0	2.5	43.5	2.8	28.4	8.6	54.5	66.9	27.8	9.5	6.2	65.0
Seedway	SW 1488TR 7364PCE	114 113	43 33	33,000 33,000	40.3 40.1	8.2 7.6	2.2	3.0 2.4	40.8 45.3	2.5	29.4 27.8	9.1 8.5	54.9 53.0	67.9 67.6	24.0 27.1	8.2 9.2	5.3 6.0	65.4 64.9
Chemgro				33,000	39.8	7.6	2.1	2.4	45.3		27.8	9.0			27.1	9.2	6.0	64.9
Kings Agriseeds	RedTail RT 64T39	114	17 19	· ·						2.8			55.1	66.6				
Augusta	A1964	114	33	33,000	39.6	7.5 7.6	2.1	2.4	44.1 42.8	2.6 2.8	28.8	8.7 8.9	54.9 55.3	68.4	29.1 24.6	10.0 8.4	6.5	65.6 65.1
Tidewater Seed	66V66	112		33,000	39.5						29.7			67.6			5.5	
Augusta	A2162	112	10	33,000	39.2	7.7 7.8	2.1	2.4	42.7	2.7	29.3	8.7	55.2 55.7	66.8	25.9	8.8	5.7 5.7	65.0
Syngenta	NK1480-DV	114	19	33,000	39.2	7.8	2.3	2.7	39.0		32.1	9.5		66.7	25.9	8.8		64.7
Mid-Atlantic	MA5144D	114	17	33,000	39.0		2.3	2.4	42.4	2.9	29.2	8.9	55.2	67.4	27.0	9.2	6.0	65.3
Mid-Atlantic	MA6120PCE	112	32	33,000	38.9	7.9	2.2	2.7	40.4	2.8	31.0	9.4	55.5	67.1	23.3	7.9	5.2	64.9
Growmark FS	INVISION FS 6157T RIB	111	43	33,000	38.8	7.5	1.9	2.3	43.7	2.8	29.6	9.0	56.4	67.6	28.0	9.6	6.3	65.7
Pine Creek Seeds	R6317DV	113	19	33,000	38.8	7.9	2.2	2.7	41.9	2.7	29.2	9.4	55.2	67.5	25.6	8.7	5.7	65.2
445 440			111-1	14 day means	40.3	7.8	2.1	2.5	43.8	2.8	28.6	8.6	55.6	67.3	27.0	9.2	6.0	65.5
115-118 day hybrids	1839 TCRIB	118	43	33.000	43.2	7.7	2.0	2.3	47.1	2.9	26.5	8.0	55.4	68.1	31.4	10.7	7.1	66.2
Revere				,	43.2	7.7	2.0		48.5		26.5	7.7	55.4	68.3	28.0	-	6.4	
Stine Seed Tidewater Seed	9818-32 64A29	117 115	20 43	33,000 33,000	41.9	7.6	1.9	2.2	46.3	2.8	26.9	7.7	58.2	67.7	27.4	9.6 9.4	6.3	66.4 66.8
Dekalb	DKC115-81RIB	115	40	33,000	41.8	7.8	2.3	2.5	42.9	2.8	29.3	9.2	53.7	66.5	25.0	8.5	5.5	64.3
				•														
Advanta Seeds	XC25231	117	19	33,000	41.7	7.6	2.1	2.1	47.0	3.0	26.7	8.6	54.1	67.2	26.8	9.2	6.0	65.2
Revere	1627 TCRIB	116	43	33,000	41.5	7.7	2.2	2.7	42.7	2.6	29.2	8.9	54.5	68.3	26.8	9.1	6.0	65.4
Pine Creek Seeds	R6516PC	115	33	33,000	41.3	7.9	2.1	2.5	45.9	2.8	26.7	8.3	55.6	67.5	26.9	9.2	6.0	65.9
Channel	215-84SSPRIB	115	40	33,000	41.1	7.8	2.0	2.6	44.2	2.7	28.7	8.2	55.7	67.4	26.3	9.0	5.9	65.5
Syngenta	E118K9-DV	118	19	33,000	40.6	7.7	2.3	2.6	43.2	2.7	28.8	9.0	52.8	67.0	24.9	8.5	5.4	64.3
Chemgro	7769SSP	117	40	33,000	40.5	7.7	2.2	2.3	44.9	2.8	28.2	8.9	53.0	66.9	27.6	9.4	6.1	64.5
Syngenta	E117Z7-D	117	18	33,000	40.4	7.6	2.0	2.6	43.2	2.6	28.6	8.7	54.7	67.9	27.7	9.4	6.2	65.5
Stine Seed	9817-30	116	45	33,000	39.5	8.1	1.9	2.6	44.3	2.8	27.4	8.7	54.5	66.6	27.6	9.4	6.1	65.0
Seedway	SW 1579SS	115	39	33,000	39.3	7.9	2.3	2.6	42.6	2.8	29.2	9.0	55.4	66.9	26.5	9.0	5.9	65.1
Channel	215-40VT4PRIB	115	53	33,000	38.8	7.4	2.4	2.4	39.3	2.5	32.3	10.0	51.8	67.4	26.5	9.1	5.7	63.3
Advanta Seeds	XC25343	117	1	33,000	37.7	7.7	2.4	2.6	39.7	2.6	31.1	10.2	53.0	68.2	24.8	8.4	5.5	64.5
Seed Consultants	SC1185V	118	48	33,000	36.2	7.6	2.3	2.6	39.4	2.5	31.6	10.0	53.1	67.2	26.4	9.0	5.8	63.9
			115-1	18 day means	40.5	7.7	2.2	2.5	43.8	2.7	28.6	8.8	54.4	67.4	26.9	9.2	6.0	65.1
				Overall Mean	40.4	7.7	2.1	2.5	43.8	2.7	28.6	8.7	55.1	67.4	27.0	9.2	6.0	65.3
				LSD(0.1)	NS	NS	0.3	NS	NS	NS	NS	NS	NS	NS	3.1	1.1	0.7	NS
				CV%	7.6	4.6	9.9	14.9	9.6	9.4	11.1	14.1	5.0	1.3	8.5	8.6	9.1	1.9
				CV%	7.0	4.0	3.3	14.5	9.0	7.4	11.1	14.1	5.0	1.5	0.5	0.0	J.1	1.5

PDMP Corn Silage Hybrid Testing Program 2025



College of Agricultural Sciences

Late maturity (111-118) day RM silage hybrids in West Grove, PA

Notes: SEE BACKGROUND TAB

Cooperator: Walmoore Holsteins Inc.

in collaboration with

PennState Extension

					Dry	Crude						uNDF			Fresh		DOM	
		Relative		Pop.	Matter	Protein	Lignin	Ash	Starch	TFA	NDFom	240 hr	NDFD 30	IVSD	Yield	OM Yield	Yield	OMD
Brand	Hybrid	Maturity	Traits ¹	Plants/ac	% ²	%DM	%DM	%DM	%DM	%DM	%DM	%DM	%NDF	%Starch ³	tons/ac⁴	tons/ac⁵	tons/ac ⁶	% ⁷

Traits: See tab " Trait Key" for individual trait designation.

⁷OMD: Organic Matter Digestibility - Please see "OMD Story" tab for information on how to use this column

NS = Not Significant

Prepared by: Alex Hristov (PSU Animal Sciences), Sergio Francisco (Cargill), Chris Canale (Cargill), Hanna Wells(PSU Plant Science), Dayton Spackman (PSU Plant Science), Cassidy Bumbaugh (PSU Plant Science), Charlie White (PSU Plant Science)

² **Dry Matter:** Tables are sorted by dry matter. <u>Avoid making comparisons with hybrids that differ significantly in dry matter.</u>

³ IVSD: Starch digestibiliy (% of starch) is analyzed by an NIRS method on samples ground through a 4-mm screen and incubated for 7 hours (IVSD).

⁴ Fresh Yield: Silage yields are expressed on a 35 percent DM basis; all other parameters are expressed on a dry matter basis.

⁵ **OM Yield:** Silage yield (tons/ac) expressed on an organic matter (OM) basis.

DOM Yield: Yield of digestible organic matter.

Compiled byChris DiFonzo
Michigan State University

Web site hosted by Pat Porter Texas A&M University

The most up-to-date version and related extension materials are free online at: www.texasinsects.org/bt-corn-trait-table.html Questions? difonzo@msu.edu

	TABLE 2 Principal trait packages available in	lottor	Traits in the package **********		Exp ne or	Refuge in northern	Weed control Trait									
# ^	the U.S. (alternate names in parentheses)	letter code	Font type denotes target: caterpillar or <i>rootworm</i>	B C W	S B	S C B	S W C B	T A W	W B C	B <i>C C</i>		states (higher in the south)	*check bag tag			
0	Conventional	С														
3	AcreMax	AM	Cry1Ab Cry1F	×	R	RL	RL	х	х	RL		R			5% blend	GLY LL
4	AcreMax Leptra	AML	Cry1Ab Cry1F Vip3A	×	х	RL	х	х	х	х	х	х			5% blend	GLY LL
6	AcreMax Xtra	AMX	Cry1Ab Cry1F Cry34/35Ab1	х	R	RL	RL	×	х	RL		R	RL	R	10% blend	GLY LL
7	AcreMax Xtreme	AMXT	Cry1Ab Cry1F Cry34/35Ab1 mCry3A	х		RL	RL	×	х	RL		R	х	R	5% blend	GLY LL
11+12	Agrisure Above (Agrisure3120EZ) AA Refuge Renew (Agrisure3120)	AA	Cry1Ab Cry1F	x	R	RL	RL	х	х	RL		R			5% blend Renew: 5%	GLY LL*
13+14	Agrisure Total (Agrisure3122EZ) AT Refuge Renew (Agrisure3122)	AT	Cry1Ab Cry1F Cry34/35Ab1 mCry3A	х	R	RL	RL	х	х	RL		R	х	R	5% blend Renew: 5%	GLY LL*
15	Agrisure Viptera 3110	3110	Cry1Ab Vip3A	×	х	RL	х	х	х	х	х	х			20%	GLY LL
16	Agrisure Viptera 3111	3111	Cry1Ab Vip3A <i>mCry3A</i>	×	х	RL	х	х	х	х	х	х	х	R	20%	GLY LL
17+18	Duracade (Agrisure5122EZ) D Refuge Renew (Agrisure5122)	D	Cry1Ab Cry1F eCry3.1Ab mCry3A	х	R	RL	RL	х	х	RL		R	х	R	5% blend Renew: 5%	GLY LL*
19+20	Duracade Viptera (Agrisure5222EZ) DV Refuge Renew (Agrisure5222)	DV	Cry1Ab Cry1F Vip3A eCry3.1Ab mCry3A	х	х	RL	х	х	х	х	х	х	х	R	5% blend Renew: 5%	GLY LL*
21+22	Duracade Viptera Z3 (Agrisure5332EZ) DVZ Refuge Renew (Agrisure5332)	DVZ	Cry1Ab Cry1A.105 Cry2Ab2 Vip3A eCry3.1Ab mCry3A	х	х	RL	х	х	х	х	х	х	х	R	5% blend Renew: 5%	GLY LL*
29	Intrasect	YHR	Cry1Ab Cry1F	×	R	RL	RL	х	х	RL		R			5%	GLY LL
30	Leptra	VYHR	Cry1Ab Cry1F Vip3A	×	х	RL	х	х	х	х	х	х			5%	GLY LL
32	PowerCore Refuge Adv.	PWRA	Cry1A.105 Cry2Ab2 Cry1F	×	R	RL	х	х	х	RL		R			5% blend	GLY LL
33	PowerCore Enlist or Enlist Refuge Advanced	PWE PCE	Cry1A.105 Cry2Ab2 Cry1F	х	R	RL	х	х	х	RL		R			5% Adv 5% blend	GLY LL Enlist
57	PowerCore Ultra Enlist or Ultra Enlist Refuge Advanced	PWUE PCUE	Cry1A.105 Cry2Ab2 Cry1F Vip3A	х	х	RL	х	х	х	х	х	х			5% Adv 5% blend	GLY LL Enlist
34	QROME	Q	Cry1Ab Cry1F Cry34/35Ab1 mCry3A	х	R	RL	RL	х	х	RL		R	х	R	5% blend	GLY LL
35	SmartStax or Genuity SS	SS SX	Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1	х	R	RL	х	х	х	RL		R	RL	R	5%	GLY LL
36	SmartStax Enlist SS Enlist Refuge Advanced	SSE	Same as SmartStax	×	R	RL	х	х	х	RL		R	RL	R	5% Adv 5% blend	GLY LL Enlist
38	SmartStax Refuge Advanced SmartStax RIB Complete	SXRA	Same as SmartStax	х	R	RL	х	х	х	RL		R	RL	R	5% blend	GLY LL
40	SmartStax PRO	SSPro	Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1 dvSnf7	х	R	RL	х	х	х	RL		R	х	х	5%	GLY LL
41	SmartStax PRO Enlist SSPro Enlist Refuge Advanced	SSPro	Same as SmartStax Pro	х	R	RL	х	х	х	RL		R	х	х	5% Adv 5% blend	GLY LL Enlist
42	SmartStax PRO Refuge Advanced RIB Complete or w/RNAi Tech	SSPro	Same as SmartStax Pro	×	R	RL	х	х	х	RL		R	х	х	5% blend	GLY LL
43	Trecepta RIB Complete	TRERIB	Cry1A.105 Cry2Ab2 Vip3A	х	х	RL	х	х	х	х	х	х			5% blend	GLY
44+45	Viptera (Agrisure3220EZ) Vip Refuge Renew (Agrisure3220)	V	Cry1Ab Cry1F Vip3A	х	х	RL	х	х	х	х	х	х			5% blend Renew: 5%	GLY LL*
46+47	Viptera Z3 (Agrisure3330EZ) VZ Refuge Renew (Agrisure3330)	VZ	Cry1Ab Cry1A.105 Cry2Ab2 Vip3A	х	х	RL	х	х	х	х	х	х			5% blend Renew: 5%	GLY LL*
48	Vorceed Enlist	V	Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1 dvSnf7	х	R	RL	х	х	х	RL		R	х	х	5% blend	GLY LL Enlist
NA	Vorceed Enlist Structured - Expected in 2026	VS	Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1 dvSnf7	х	R	RL	х	х	х	RL		R	х	х	5%	GLY LL Enlist
49	VT Double PRO	VT2P	Cry1A.105 Cry2Ab2		R	RL	х	х	х	RL					5%	GLY
50	VT2 PRO RIB Complete	VT2PRIB	Cry1A.105 Cry2Ab2		R	RL	х	х	х	RL					5% blend	GLY
52	VT3 PRO RIB Complete	VT3PRIB	Cry1A.105 Cry2Ab2 Cry3Bb1		R	RL	х	х	х	RL			RL	R	10% blend	GLY
53	VT4 PRO w/RNAi Technology	VT4PRO	Cry1A.105 Cry2Ab2 Vip3A Cry3Bb1 dvSnf7	х	х	RL	х	х	х	Х	х	х	Х	х	5% blend	GLY

The OMD Index

The digestibility of nutrients in corn silage is paramount when determining nutritional value. Starch and NDF are responsible for much of the digestible energy in corn silage. In order to give dairy producers and nutritionist a tool to evaluate corn silage hybrids, we developed a new digestibility index, called the Organic Matter Digestibility Index (OMDI or just OMD), and is based on digestibility of protein, fat, NDF, and starch. The sum of which makes up approximately 86-88% of the organic matter in corn silage.

The OMD index represents the digestible portion of silage organic matter and is based on chemical analyses only. It does not predict dry matter intake or milk production, although numerous studies clearly show that digestibility of forage organic matter is directly related to lactation performance of dairy cows. The OMD index does not represent the absolute digestibility of silage organic matter, as this can be reliably determined only in experiments with live animals. But, OMD is representative of the potentially digestible organic matter of the whole plant and can be used to compare silage hybrids. Furthermore, simulation analyses using the Cornell Net Carbohydrate and Protein System (CNCPS v. 6.55; Cornell University, Ithaca, NY) show that OMD correlates reasonably well with model-predicted milk production of dairy cows fed a standard diet containing approx. 40% corn silage (dry matter basis).

How is the OMD Index Used?

Feeding value of corn silage is mostly associated with digestibility of NDF or starch. A long-standing goal of PDMP is to create a single measure of silage nutritive value using several variables associated with digestibility. Traditional variables, crude protein (accounted for fiber-bound nitrogen), NDF, starch, lignin, and fat, are combined with digestibility determinations for NDF (NDFD30*) and starch (IVSD; 7-hour, 1-mm grind). Once combined, these digestibility coefficients sum to predict OMD.

The OMD Index is calculated using the following equation: OMDI (%) = $\{[(\text{crude protein} - \text{NDICP}) \times 0.89] + (\text{total fatty acids} \times 0.75) + (\text{starch} \times \text{IVSD} \div 100) + [(\text{aNDFom} - \text{lignin}) \times \text{NDFD30} \div 100)]\} \div [(\text{crude protein} - \text{NDICP}) + \text{total fatty acids} + \text{starch} + (\text{aNDFom} - \text{lignin})] \times 100.$

Where: OMDI (%) is Organic Matter Digestibility Index; crude protein, total fatty acids, starch, NDICP (NDF-bound crude protein), aNDFom (ash-free basis, amylase-treated NDF), and lignin (ash-free) are expressed as % of corn silage dry matter; 0.89 is assumed (based on literature data) coefficient of digestibility of silage crude protein; 0.75 is assumed (based on literature data) coefficient of digestibility of silage total fatty acids; IVSD is starch digestibility (by NIRS at 7-hour and sample ground through a 4-mm sieve) expressed as % of starch; and NDFD30.

Use of OMDI: The OMD index is intended to represent the digestible portion of silage dry matter and is based on chemical analyses. OMD does not represent the absolute digestibility of silage organic matter, but it is representative of the potentially digestible organic matter and can be used when comparing silage hybrids. **Simply put, the higher the OMD value, the higher the overall expected digestibility of the silage.** OMD reflects the digestibility of key nutrients within the entire plant. Producers without carryover of silage should consider the interaction of OMD and DOM (digestible organic matter yield per acre) as yield of digestible organic matter will be equally as relevant as OMD.

Conclusion

Organic matter digestibility is not a new measure. For years, researchers and nutritionists have used digestibility estimates to formulate rations for dairy cattle. Today, integrating these data is a useful practice to gauge silage value and match hybrid to farm needs. Put simply, OMD measures whole plant digestibility. Emphasis is on digestibility of all main nutrients. In the end, we hope OMD serves to facilitate discussion among producer, seed consultant, and dairy nutritionist as to which hybrids offer the best nutrient value for dairy cows.